References:

- 1. Kempthorne P, Morriss WW, Mellin-Olsen J, Gore-Booth J. The WFSA Global Anesthesia Workforce Survey. Anesthesia & Analgesia 2017; **125**: 981-90.
- 2. Jenkins K, Wong DT, Correa R. Management choices for the difficult airway by anesthesiologists in Canada. Canadian Journal of Anesthesia 2002; **49**: 850-6.
- 3. Rosenblatt WH, Wagner PJ, Ovassapian A, Kain ZN. Practice Patterns in Managing the Difficult Airway by Anesthesiologists in the United States. Anesthesia & Analgesia 1998; **87**: 153-7.
- 4. Kreijchie RV, Morgan DW. Determining Sample Size for Research Activities. Educational and Psychological Measurement 1970; **30**: 607-10.
- 5. Holmes MG, Dagal A, Feinstein BA, Joffe AM. Airway Management Practice in Adults With an Unstable Cervical Spine: The Harborview Medical Center Experience. Anesthesia And Analgesia 2018; **127**: 450-4.
- 6. Lord SA, Boswell WC, Williams JS, Odom JW, Boyd CR. Airway control in trauma patients with cervical spine fractures. Prehospital And Disaster Medicine 1994; **9**: 44-9.
- 7. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. Journal of Biomedical Informatics. 2009;42(2):377-81.